

NNLO corrections to $\bar{B} \rightarrow X_u \ell \bar{\nu}$ and the determination of $|V_{ub}|$

Ben D. Pecjak

Institut für Physik (THEP), Johannes Gutenberg-Universität, D-55099 Mainz, Germany

The calculation of partial decay rates in $\bar{B} \rightarrow X_u \ell \bar{\nu}$ decays at next-to-next-to-leading order (NNLO) in α_s and to leading order in $1/m_b$ is described. New results for the hard function are combined with known results for the jet function and shape-function moments in a numerical analysis which explores the impact of the NNLO corrections on partial decay rates and the determination of $|V_{ub}|$.

1. INTRODUCTION

The inclusive decay $\bar{B} \rightarrow X_u \ell \bar{\nu}$ is of much interest because of its potential to constrain the CKM element $|V_{ub}|$. Due to experimental cuts required to suppress charm background, measurements of this decay are available only in the shape-function region, where the hadronic final state is collimated into a single jet carrying a large energy on the order of m_b and a moderate invariant mass squared on the order of $m_b \Lambda_{\text{QCD}}$. The theory challenge is to calculate partial decay rates in the presence of these cuts, where a local operator product expansion is insufficient. A systematic treatment relies on a non-local operator product expansion whose end result can be formulated in terms of a factorization theorem. Different approaches to this factorization have been put forth in the literature, going under the names of BLP [1, 2], GGOU [3], and the dressed-gluon exponentiation [4]. The purpose of this talk is to describe the elements that go into the BLP formalism at next-to-next-to-leading order (NNLO) in α_s and leading order in $1/m_b$.

The main point of discussion is the factorization formula for an arbitrary decay distribution restricted to the kinematics of the shape-function region

$$d\Gamma \sim H \cdot J \otimes S + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}}{m_b}\right), \quad (1)$$

where the symbol $\otimes$ denotes a convolution. The factorization formula contains a hard function H , which is related to physics at the hard scale m_b , a jet function J , which is related to physics at the intermediate scale $m_b \Lambda_{\text{QCD}}$, and a non-perturbative shape function S , describing the internal soft dynamics of the B meson [5, 6]. This factorized form was originally derived in [7, 8] using diagrammatic techniques, and was rederived in the framework of soft-collinear effective theory (SCET) in [1, 9]. The hard and jet functions to NLO in perturbation theory have been known for some time [1, 9], and the NNLO jet function was obtained in [10]. Very recently, the NNLO contributions to the hard function have also been calculated [11, 12, 13], thus completing the perturbative corrections to the factorization formula to this order. In Section 2 we review the calculation of H and J to NNLO, along with the form and solutions of their renormalization-group (RG) evolution equations. Then, in Section 3, we give preliminary results illustrating the numerical impact of the NNLO corrections on the P_+ spectrum, comment on the relevance for the determination of $|V_{ub}|$, and make some concluding remarks.

2. THE HARD AND JET FUNCTIONS AT NNLO

In this section we describe the calculation of the hard and jet functions at NNLO, as well as the form and solution of their RG evolution equations.

The hard function arises when integrating out fluctuations at the scale m_b by matching QCD onto SCET. This matching can be done at the level of the $b \rightarrow u$ transition current, and to leading order in $1/m_b$ takes the form

$$e^{-im_b v \cdot x} \bar{u}(x) \gamma^\mu (1 - \gamma_5) b(x) = \sum_{i=1}^3 \int ds \tilde{C}_i(s) \bar{\chi}(x + s\bar{n}) \Gamma_i^\mu \mathcal{H}(x_-), \quad (2)$$

where we have followed the SCET conventions of [1], and the Γ_i^μ are a set of three Dirac structures. In practice, the matching calculation is carried out in momentum space and yields results for the Fourier-transformed coefficients, which read

$$C_i(\bar{n} \cdot p) = \int ds e^{is\bar{n} \cdot p} \tilde{C}_i(s). \quad (3)$$

The hard function H is derived from the matrix of coefficients $H_{ij} = C_i C_j$.

To obtain the matching coefficients C_i requires to calculate UV-renormalized matrix elements in full QCD and SCET. The calculation is simplest when the external states are chosen as on-shell quarks and both UV and IR divergences are regulated in dimensional regularization in $d = 4 - 2\epsilon$ dimensions. In that case the loop corrections to the SCET matrix elements are given by scaleless integrals and vanish, so that the result is just its tree-level value multiplied by renormalization factors from operator and wave-function renormalization. The main challenge is to calculate the QCD result, which is written in terms of three Dirac structures multiplied by scalar form factors D_i . To obtain these scalar form factors at NNLO requires to calculate the two-loop corrections to the $b \rightarrow u$ current in QCD. This task has been completed in [11, 12, 13]. One then turns the results for the scalar form factors D_i into those for the SCET Wilson coefficients by evaluating the matching conditions

$$C_i(m_b, \bar{n} \cdot p, \mu) = \lim_{\epsilon \rightarrow 0} Z_J^{-1}(\epsilon, m_b, \bar{n} \cdot p, \mu) D_i(\epsilon, m_b, \bar{n} \cdot p, \mu), \quad (4)$$

where Z_J is a current renormalization factor. Details of the matching calculation and results for the Wilson coefficients have been presented in [12, 13].

In the BLNP approach, the hard function is evaluated at a scale $\mu_i \sim 1.5$ GeV and logarithms of the ratio μ_h/μ_i are treated as large. One can resum these logarithms by deriving and solving the RG equation for the hard coefficients C_i , or equivalently the matrix of coefficients H_{ij} . The evolution equation has the form (see, e.g. [1])

$$\frac{d}{d \ln \mu} H_{ij}(\bar{n} \cdot p, \mu) = 2 \left[\gamma'(\alpha_s) + \Gamma_{\text{cusp}}(\alpha_s) \ln \frac{\bar{n} \cdot p}{\mu} \right] H_{ij}(\bar{n} \cdot p, \mu), \quad (5)$$

and its solution is

$$H_{ij}(\bar{n} \cdot p, \mu) = y^{-2a_\Gamma(\mu_h, \mu)} \exp \left[2S(\mu_h, \mu) - 2a_\Gamma(\mu_h, \mu) \ln \frac{m_b}{\mu_h} - 2a_{\gamma'}(\mu_h, \mu) \right] H_{ij}(\bar{n} \cdot p, \mu_h), \quad (6)$$

where we have defined $y = \bar{n} \cdot p/m_b$. Explicit results for the Sudakov factor S and the anomalous exponents a_Γ , $a_{\gamma'}$ can be read off from [1]. A consistent treatment at NNLO in α_s requires the matching coefficients at two loops, the anomalous dimension γ' at three loops, and the cusp anomalous dimension at four loops. The anomalous dimensions are both known to one loop lower, which adds a small uncertainty to the analysis.

We now turn to the jet function J , which arises when integrating out fluctuations at the intermediate scale $m_b \Lambda_{\text{QCD}}$ by matching SCET onto HQET. The calculation of the jet function can be recast into the evaluation of the imaginary part of a certain vacuum matrix element in QCD, and was calculated to NNLO in [10]. In light-cone gauge, this matrix element is the quark propagator. The jet function depends on a renormalization scale μ , which is usually kept fixed at $\mu_i = 1.5$ GeV in the BLNP approach. However, it is possible to derive and solve the RG equation for J , which allows one to also vary the renormalization scale at which the jet function is evaluated. The starting point is the exact integro-differential evolution equation derived in [10]:

$$\frac{dJ(p^2, \mu)}{d \ln \mu} = - \left[2\Gamma_{\text{cusp}} \ln \frac{p^2}{\mu^2} + 2\gamma^J \right] J(p^2, \mu) - 2\Gamma_{\text{cusp}} \int_0^{p^2} dp'^2 \frac{J(p'^2, \mu) - J(p^2, \mu)}{p^2 - p'^2}. \quad (7)$$

A novel solution to the evolution equation (7) was presented in [15]. The method is based on the observation that the Laplace transformed jet function

$$\tilde{J}\left(\ln \frac{Q^2}{\mu^2}, \mu\right) = \int_0^\infty dp^2 e^{-sp^2} J(p^2, \mu), \quad s = \frac{1}{e^{\gamma_E} Q^2} \quad (8)$$




Figure 1: Partial rates with a cut $P_+ < M_D^2/M_B$ as a function of μ_h at LO, NLO, and NNLO.

obeys a local evolution equation analogous to (5). Solving as before and inverting the Laplace transform gives the resummed J in momentum space. Alternatively, it is always possible to write partial decay rates in terms of an integral over the jet function, defined as

$$j\left(\ln \frac{Q^2}{\mu^2}, \mu\right) = \int_0^{Q^2} dp^2 J(p^2, \mu). \quad (9)$$

Using the solution of the RG equation, one can write a compact expression for the resummed integrated jet function

$$j\left(\ln \frac{Q^2}{\mu_f^2}, \mu_f\right) = \exp[-4S(\mu_i, \mu_f) + 2a_{\gamma J}(\mu_i, \mu_f)] \left(\frac{Q^2}{\mu_i^2}\right)^\eta \tilde{j}\left(\partial_\eta + \ln \frac{Q^2}{\mu_i^2}, \mu_i\right) \frac{e^{-\gamma_E \eta}}{\Gamma(1+\eta)}, \quad (10)$$

where expressions for the RG factors $a_{\gamma J}$ and η can be found in [15]. Equations (6) and (10) allow one to evaluate integrals of the product $H \cdot J$ at an arbitrary scale μ_f at which the shape function is renormalized, and study the dependence of partial decay rates under independent variations of the matching scales μ_h and μ_i , under which they are formally independent. Detailed numerical results shall be given in [14].

3. NUMERICAL RESULTS AND CONCLUSIONS

In this section we briefly explore the numerical impact of the NNLO corrections, taking as an example the partial rate with a cut $P_+ < \Delta = M_D^2/M_B$. This partial rate can be written as

$$\Gamma_u(P_+ < \Delta) = \frac{G_F^2 |V_{ub}|^2}{96\pi^3} \int_0^\Delta d\hat{\omega} \hat{S}(\hat{\omega}, \mu_f) \int_0^{M_B - \hat{\omega}} dq 5q^4 \int_0^1 dy y^{2-2a\Gamma} H_u(y, \mu_f) j\left(\ln \frac{m_b y(\Delta_q - \hat{\omega})}{\mu_f}, \mu_f\right) \quad (11)$$

where $\Delta_q = \min(\Delta, M_B - q)$, and the hard and integrated jet functions are to be evaluated in their resummed forms (6) and (10). The function H_u is related to certain combinations of the H_{ij} and can be deduced from [2]. To evaluate (11) requires a model for the shape function. In what follows, we use the two-parameter exponential model

$$\hat{S}(\hat{\omega}) = \mathcal{N}(b, \Lambda) \hat{\omega}^{b-1} \exp\left(-\frac{b\hat{\omega}}{\Lambda}\right). \quad (12)$$

It is possible to put model-independent constraints on the normalization factor $\mathcal{N}$ and the parameters b and Λ by studying shape-function moments [1]. These moments are defined as

$$M_N(\hat{\omega}_0, \mu_i) \equiv \int_0^{\hat{\omega}_0} d\hat{\omega} \hat{\omega}^N \hat{S}(\hat{\omega}, \mu_i), \quad (13)$$

and can be obtained as an expansion in local HQET operators as long as $\hat{\omega}_0 \gg \Lambda_{\text{QCD}}$. The zeroth moment fixes the normalization of the shape function and was determined to NNLO in [16], and can be used along with the first and

second moment to define the parameters $m_b = M_B - \bar{\Lambda}$ and μ_π^2 in the shape-function scheme. Numerical values for the HQET parameters m_b and μ_π^2 in the shape-function scheme at NNLO can be determined from those obtained from global fits in other schemes using the two-loop conversion relations derived in [17]. One then tunes the parameters b and Λ such that the moments of the model function (12) give appropriate results for m_b and μ_π^2 . In our analysis here, we tune the parameters such that $m_b = 4.61$ GeV and $\mu_\pi^2 = 0.2 \text{ GeV}^2$ at NNLO in the shape-function scheme.

In Figure 1, we show results for the partial rate with a cut $P_+ < \Delta = M_D^2/M_B$, for the choice $\mu_i = \mu_f = 1.5$ GeV, as a function of the hard matching scale μ_h . From the figure, one sees that the dependence on the renormalization scale μ_h is reduced when going to higher orders in perturbation theory, and also that there is a fairly large downward shift in the central value between NLO and NNLO. This would tend to raise the value of $|V_{ub}|$ compared to the result deduced from the NLO calculation in the BLP approach. However, we must stress that these numerical results are preliminary, and will be finalized in [14], along with uncertainties associated with variations of the matching scale μ_i . Results for other partial rates, such as those with cuts on the hadronic invariant mass and lepton energy, will also be presented.

It is worth emphasizing that the partial decay rates are rather sensitive to the numerical value of m_b . If, as suggested in [18], the $B \rightarrow X_s \gamma$ moments are excluded from global fits on the grounds that measurements of the photon energy spectrum are made in kinematic regions where shape-function effects are important, then m_b tends to come out closer to 4.71 GeV. Changing m_b by such an amount raises the partial decay rates upwards by approximately 20%. A precise determination of $|V_{ub}|$ from inclusive decays will thus require that this point be settled.

Acknowledgments

I am grateful to Hrachia Asatryan, Christoph Greub, and Matthias Neubert, for collaborations on the work which formed the basis of this talk.

References

- [1] S. W. Bosch, B. O. Lange, M. Neubert and G. Paz, Nucl. Phys. B **699** (2004) 335 [arXiv:hep-ph/0402094].
- [2] B. O. Lange, M. Neubert and G. Paz, Phys. Rev. D **72** (2005) 073006 [arXiv:hep-ph/0504071].
- [3] P. Gambino, P. Giordano, G. Ossola and N. Uraltsev, JHEP **0710** (2007) 058 [arXiv:0707.2493 [hep-ph]].
- [4] J. R. Andersen and E. Gardi, JHEP **0601** (2006) 097 [arXiv:hep-ph/0509360].
- [5] M. Neubert, Phys. Rev. D **49**, 3392 (1994) [arXiv:hep-ph/9311325]; M. Neubert, Phys. Rev. D **49**, 4623 (1994) [arXiv:hep-ph/9312311].
- [6] I. I. Y. Bigi, M. A. Shifman, N. G. Uraltsev and A. I. Vainshtein, Int. J. Mod. Phys. A **9**, 2467 (1994) [arXiv:hep-ph/9312359].
- [7] G. P. Korchemsky and G. Sterman, Phys. Lett. B **340**, 96 (1994) [arXiv:hep-ph/9407344].
- [8] R. Akhouchy and I. Z. Rothstein, Phys. Rev. D **54** (1996) 2349 [arXiv:hep-ph/9512303].
- [9] C. W. Bauer and A. V. Manohar, Phys. Rev. D **70** (2004) 034024 [arXiv:hep-ph/0312109].
- [10] T. Becher and M. Neubert, Phys. Lett. B **637**, 251 (2006) [arXiv:hep-ph/0603140].
- [11] R. Bonciani and A. Ferroglia, arXiv:0809.4687 [hep-ph].
- [12] H. M. Asatryan, C. Greub and B. D. Pecjak, arXiv:0810.0987 [hep-ph].
- [13] M. Beneke, T. Huber and X. Q. Li, arXiv:0810.1230 [hep-ph].
- [14] H. M. Asatryan, C. Greub, M. Neubert, and B. D. Pecjak, in preparation.
- [15] T. Becher and M. Neubert, Phys. Rev. Lett. **97** (2006) 082001 [arXiv:hep-ph/0605050].
- [16] T. Becher and M. Neubert, Phys. Lett. B **633**, 739 (2006) [arXiv:hep-ph/0512208].
- [17] M. Neubert, Phys. Lett. B **612**, 13 (2005) [arXiv:hep-ph/0412241].
- [18] M. Neubert, arXiv:0801.0675 [hep-ph].